

Tilburg University

Parallel normreducing transformations for the algebraic eigenvalue problem

Paardekooper, M.H.C.

Publication date:
1990

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Paardekooper, M. H. C. (1990). *Parallel normreducing transformations for the algebraic eigenvalue problem*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 423). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
CBM
R



— KE
UNIVERSITEIT
BRABANT

7626
1990
423

3

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS



* C I N O 1 1 9 8 *



DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM



**PARALLEL NORMREDUCING TRANSFORMATIONS
FOR THE ALGEBRAIC EIGENVALUE PROBLEM**

M.H.C. Paardekooper

R 83

FEW 423

518.315

PARALLEL NORMREDUCING TRANSFORMATIONS
FOR THE ALGEBRAIC EIGENVALUE PROBLEM

M.H.C. Paardekooper

Department of Econometrics
Tilburg University
5000 LE Tilburg

This research is part of the VF-program "Parallele Algoritmiëk", TD-WI-08185-25, which has been approved by the Netherlands Ministry of Education and Sciences.

PARALLEL NORMREDUCING TRANSFORMATIONS FOR THE ALGEBRAIC EIGENVALUE PROBLEM

ABSTRACT

This article presents a unified approach for parallel normreducing methods for the algebraic eigenproblem. The so-called Euclidean parameters presents the problem, to minimize the Frobenius norm of the transform matrix, in a simple form. The use of appropriate preprocessing unitary transforms together with an appropriate pivot strategy leads to convergence to normality.

Keywords: Jacobi methods, parallel transformations, eigenvalues, convergence to normality, Euclidean parameters, normreduction, commutator.

INTRODUCTION

In 1971 Sameh [5] proposed a Jacobi-like eigenvalue algorithm for a parallel computer. Sameh's method is a parallel elaboration of Eberlein's sequentially normreducing transformation procedure [1,4].

In this paper we present a unified approach to Jacobi-like normreducing transformations and we apply it to elucidate and improve Sameh's method. In a sequentially Jacobi-like procedure for the computation of the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ of a real or complex matrix $A = A^{(0)}$ (order n) a sequence $\{A^{(j)}\}$ is constructed recursively:

$$A^{(j+1)} = T^{(j)-1} A^{(j)} T^{(j)}, \quad j \geq 0. \quad (1.1)$$

In (1.1) $T^{(j)}$ is a unimodular shear matrix with Jacobi parameter $(p_j, q_j, r_j, s_j) \in \mathbb{C}^4$ in (ℓ, m) -restriction \hat{T}_{ℓ_j, m_j} of $T_{\ell_j, m_j} = T^{(j)}$

$$\hat{T}_{\ell_j, m_j} = \begin{bmatrix} p_j & q_j \\ r_j & s_j \end{bmatrix} \begin{matrix} \leftarrow \ell_j \\ \leftarrow m_j \end{matrix} \quad \begin{matrix} \uparrow \\ \uparrow \end{matrix} \begin{matrix} \ell_j \\ m_j \end{matrix} \quad (1.2)$$

In Eberlein-like normreducing processes [1,3,6] the aim is to construct $\{A^{(j)}\}$ so that

$$\lim_{k \rightarrow \infty} \|A^{(j)}\|_F = \sum_{j=1}^n |\lambda_j|^2, \quad (1.3)$$

that means $\{A^{(j)}\}$ converges to normality [2]. Eberlein [1] gives for each iteration an approximation of the optimal normreducing $T_{\ell, m}$. These choices of the Jacobi parameters, together with a well-defined pivot strategy $\{(\ell_j, m_j)\}$ brings about convergence to normality.

Since the Frobenius norm is invariant under unitary transformations, the optimal norm-reducing shear $T_{\ell, m}$ is determined except for a unitary factor. Hence we discuss the normreduction in theoretical terms that are invariant under unitary transformations.

Matrices $S, P \in \mathbb{C}^{n \times n}$ will be called *row-congruent* if and only if $S = PU$ for some unitary U , notation $S \sim P$. It is easy to see that $S \sim P$ if and only if $SS^* = PP^*$. Now for shear $T_{\ell, m}$, with $\hat{T}_{\ell, m} = \begin{bmatrix} p & q \\ r & s \end{bmatrix}$

$$\hat{T}_{\ell, m} \hat{T}_{\ell, m}^* = \begin{bmatrix} |p|^2 + |q|^2 & p\bar{r} + q\bar{s} \\ \bar{p}r + \bar{q}s & |r|^2 + |s|^2 \end{bmatrix} = \begin{bmatrix} x & z \\ \bar{z} & y \end{bmatrix} \quad (1.3)$$

The quantities

$$\begin{aligned} x = x(T_{\ell, m}) &= |p|^2 + |q|^2, \quad y = y(T_{\ell, m}) = |r|^2 + |s|^2, \\ z = z(T_{\ell, m}) &= u + iv = p\bar{r} + q\bar{s} \end{aligned} \quad (1.4)$$

will be called the *Euclidean parameters* of $T_{\ell, m}$ [3]. These parameters are pre-eminent appropriate for the formulation of the Frobenius norm of $T^{-1}AT$.

We assume $T_{\ell, m}$ to be unimodular, so

$$x, y > 0 \text{ and } |ps - qr|^2 = xy - |z|^2 = 1 \quad (1.4)$$

With $z = u + iv$

$$\mathcal{H} = \{(x, y, u, v) \in \mathbb{R}^4 \mid xy - u^2 - v^2 = 1, x, y > 0\} \quad (1.5)$$

is the *positive sheet* of the elliptic hyperboloid $xy - u^2 - v^2 = 1$. In case of real norm-reducing shears, $\hat{T}_{\ell, m} \in \mathbb{R}^{2 \times 2}$,

$$x = p^2 + q^2, \quad y = r^2 + s^2, \quad z = pr + qs, \quad xy - z^2 = 1. \quad (1.6)$$

The Euclidean parameters x, y and z of that unimodular shear $T_{\ell, m}$ correspond with the points in the positive sheet of $xy - z^2 = 1$ in \mathbb{R}^3 .

In the *parallel normreduction* $A^{(j)}$ is transformed by a direct sum of *identical unimodular 2×2 matrices*:

$$A^{(j+1)} = W_j^{-1} A^{(j)} W_j, \quad j \geq 0, \quad (1.7)$$

where $W_j = \text{diag}(T_{1,j}, \dots, T_{k,j})$, $n = 2k$ and

$$T = T_{i,j} = \begin{bmatrix} p & q \\ r & s \end{bmatrix}, \quad i = 1, \dots, k. \quad (1.8)$$

Then $\|A^{(j+1)}\|_F^2$ is a quadratic function of the Euclidean parameters x, y, u and v ($z = v + iv$) of the k matrices $T_{i,j}$ in W_j . The minimization of that quartic on \mathcal{H} leads to a generalized eigenvalue problem in four dimensions. Section 2 describes the first step $A^{(1)} = W^{-1}AW$ of the normreducing process for real matrices, and there is shown the commutator $(c_{i,j}^{(1)}) = C^{(1)} = A^{(1)\top} A^{(1)} - A^{(1)} A^{(1)\top}$ in relation to parallel shear transformations: $c_{2\ell-1,2\ell}^{(1)} = 0$ and $c_{2\ell-1,2\ell-1}^{(1)} = c_{2\ell,2\ell}^{(1)}$, $\ell = 1, \dots, k$ iff transformation W minimizes $\|A^{(1)}\|_F$. This section gives also the construction of the optimal W . A special analysis is given to the step in which $\inf\{\|W_j^{-1}A^{(j)}W_j\|_F | W = T_{1,j} \otimes \dots \otimes T_{k,j}\}$ is not assumed for unimodular 2×2 shear $T_{i,j}$. Section 3 describes the same problems for complex matrices. In section 4 will be shown that a well-chosen pivot strategy $\{(\ell_{i,j}, m_{i,j}) | i = 1, \dots, k\}_{j=0}^{\infty}$ together with an appropriate preprocessing sequence of unitary matrices $U^{(j)}$ results in sequence $\{A^{(j)}\}$ that converges to normality.

2. PARALLEL NORMREDUCTION: REAL MATRICES

Let the matrix S be real and of even order $n = 2k$. Then it can be partitioned as follows

$$A = \begin{bmatrix} A_{11} & A_{12} & \dots & A_{1k} \\ A_{21} & A_{22} & \dots & A_{2k} \\ \vdots & & & \\ A_{k1} & A_{k2} & \dots & A_{kk} \end{bmatrix} \quad (2.1)$$

where each submatrix is given by

$$A_{l,m} = \begin{bmatrix} a_{2l-1,2m-1} & a_{2l-1,2m} \\ a_{2l,2m-1} & a_{2l,2m} \end{bmatrix}, \quad l, m = 1, \dots, k. \quad (2.2)$$

For convenience define

$$\begin{aligned} \sigma_{l,m} &= a_{2l,2m-1}, \quad \mu_{l,m} = a_{2l-1,2m}, \quad \alpha_{l,m} = a_{2l-1,2m-1}, \\ \beta_{l,m} &= a_{2l,2m}, \quad \nu_{l,m} = \alpha_{l,m} - \beta_{l,m}, \quad l, m = 1, \dots, k. \end{aligned} \quad (2.3)$$

Let

$$A' = W^{-1}AW, \quad (2.4)$$

where

$$W = \text{diag}(S_1, S_2, \dots, S_k)$$

with

$$S_j = \begin{bmatrix} p_j & q_j \\ r_j & s_j \end{bmatrix}, \quad p_j s_j - q_j r_j = 1, \quad j = 1, \dots, k.$$

We introduce

$$x_j = p_j^2 + q_j^2, \quad y_j = x_j^2 + s_j^2, \quad z_j = p_j r_j + q_j s_j, \quad j = 1, \dots, k;$$

they are the Euclidean parameters of the unimodular S_j . So

$$x_j y_j - z_j^2 = 1, \quad i = 1, \dots, k.$$

Then

$$A'_{\ell, m} = S_{\ell}^{-1} A_{\ell, m} S_m.$$

Easy calculations give the following results.

THEOREM 2.1. For each (ℓ, m) , $\ell, m = 1, \dots, k$, $\|A'_{\ell, m}\|_F^2$ is a bilinear function of the Euclidean parameters $(x_{\ell}, y_{\ell}, z_{\ell})$ of S_{ℓ} and (x_m, y_m, z_m) of S_m :

$$\|A'_{\ell, m}\|_F^2 = (x_{\ell}, y_{\ell}, z_{\ell}) \begin{bmatrix} \sigma_{\ell, m}^2 & \beta_{\ell, m}^2 & 2\sigma_{\ell, m} \beta_{\ell, m} \\ \alpha_{\ell, m}^2 & \mu_{\ell, m}^2 & 2\alpha_{\ell, m} \mu_{\ell, m} \\ -2\alpha_{\ell, m} \sigma_{\ell, m} & -2\beta_{\ell, m} \mu_{\ell, m} & -2(\alpha_{\ell, m} \beta_{\ell, m} + \sigma_{\ell, m} \mu_{\ell, m}) \end{bmatrix} \begin{bmatrix} x_m \\ y_m \\ z_m \end{bmatrix}.$$

THEOREM 2.2. For each $j = 1, \dots, k$ let be

$$w_j = (x_j - y_j)/2, \quad t_j = (x_j + y_j)/2 = (1 + w_j^2 + z_j^2)^{\frac{1}{2}}. \quad (2.5)$$

Then $\|W^{-1} A W\|_F^2$ is a quadratic function $(w, z) \mapsto g(w, z; A)$ where $w = (w_1, \dots, w_k)^T$ and $z = (z_1, \dots, z_k)^T$. Moreover

$$\frac{\partial g}{\partial w_{\ell}}(0, 0; A) = c_{2\ell-1, 2\ell+1} - c_{2\ell, 2\ell}, \quad \frac{\partial g}{\partial z_{\ell}}(0, 0; A) = c_{2\ell-1, 2\ell},$$

where $(c_{i, j}) = A^T A - A A^T$. \square

The complexity of the unconstrained minimization of g forces the restriction to a problem with fewer degrees of freedom. Therefore, we consider, as in [5]

$$W = \text{diag}(S_1, \dots, S_k) \quad (2.6)$$

with

$$S_j = S = \begin{pmatrix} p & q \\ r & s \end{pmatrix}, \quad ps - qr = 1, \quad j = 1, \dots, k. \quad (2.7)$$

Such a matrix W will be called a *diagonal of shears*. Now

$$x = p^2 + q^2, \quad y = r^2 + s^2, \quad z = pr - qs. \quad (2.8)$$

The unimodularity of S implies

$$xy - z^2 = 1. \quad (2.9)$$

THEOREM 2.3. If W is a diagonal of shears with Euclidean parameters (x, y, z) then

$$\|W^{-1}AW\|_F^2 = x + \sum_{\ell, m=1}^k (-\sigma_{\ell, m} x + \mu_{\ell, m} y + \nu_{\ell, m} z)^2, \quad (2.10)$$

$$\text{where } x = \sum_{\ell, m=1}^k (\text{tr}(A_{\ell, m})^2 - 2\det(A_{\ell, m})). \quad \square$$

As in (2.5) we define

$$w := (x-y)/2, \quad t := (x+y)/2 = (1+w^2+z^2)^{\frac{1}{2}}.$$

Then, as follows from (2.10), $\|W^{-1}AW\|_F^2$ is a function of w and z :

$$g(w, z; A) := \|W^{-1}AW\|_F^2. \quad (2.11)$$

With simple but cumbersome calculations one proves the following lemma.

LEMMA 2.4. Let be $(c'_{i, j}) = (A')^T A' - A'(A')^T$, where $A' = W^{-1}AW$ with W as defined in (2.6) and (2.7). Then

$$\sum_{\ell=1}^k \begin{bmatrix} c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell} \\ 2c'_{2\ell-1,2\ell} \end{bmatrix} = \begin{bmatrix} (p^2+s^2-q^2-r^2)/2 & pr-qs \\ pq-rs & ps+qr \end{bmatrix} \begin{bmatrix} \frac{\partial g}{\partial w}(w,z;A) \\ \frac{\partial g}{\partial z}(w,z;A) \end{bmatrix}, \quad (2.12)$$

where g as defined in (2.11). \square

THEOREM 2.5. The function g is stationary in $(w,z) \in \mathbb{R}^2$ iff

$$\sum_{\ell=1}^k (c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell}) = 0 \text{ and } \sum_{\ell=1}^k (c'_{2\ell-1,2\ell}) = 0.$$

PROOF. The determinant of the coefficientmatrix in (2.11) equals $\frac{1}{2}(x+y)(ps-qr) \neq 0$. \square

Theorem 2.3 implies that the determination of the optimale normreducing diagonal of shears requires the minimization of a quadratic function subject to $xy - z^2 = 1$. Let be

$$\mathbf{d} = (d_1, d_2, d_3)^T = (x, y, z),$$

$$H := \begin{bmatrix} 0 & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}. \quad (2.13)$$

and

$$\mathcal{H} := \{\mathbf{d} \in \mathbb{R}^3 \mid \mathbf{d}^T H \mathbf{d} = 1, d_1 > 0\}. \quad (2.14)$$

Further

$$B = (b_1, b_2, b_3) \in \mathbb{R}^{k^2 \times 3}$$

with

$$\mathbf{b}_1 = (\sigma_{1,1}, \sigma_{1,2}, \dots, \sigma_{k,k})^T \in \mathbb{R}^{k^2}$$

$$\mathbf{b}_2 = (\mu_{1,1}, \mu_{1,2}, \dots, \mu_{k,k})^T \in \mathbb{R}^{k^2}$$

$$\mathbf{b}_3 = (\nu_{1,1}, \nu_{1,2}, \dots, \nu_{k,k})^T \in \mathbb{R}^{k^2}$$

The Euclidean parameters $\mathbf{d} = (x, y, z)$ of an optimal normreducing diagonal of shears solve the problem

$$\min\{\|\mathbf{B}\mathbf{d}\| \mid \mathbf{d}^T \mathbf{H} \mathbf{d} = 1\}. \quad (2.15)$$

(2.15) leads to a generalized eigenproblem in three dimensions. The analysis for the three cases $\text{rank}(\mathbf{B})$ equals 3, 2 or 1 is summarized in the following theorems.

THEOREM 2.6. Let $\text{rank}(\mathbf{B}) = 3$, and $\mathbf{B} = \mathbf{Q}\mathbf{R}$ with $\mathbf{Q} \in \mathbb{R}^{k^2 \times 3}$ orthogonal and $\mathbf{R} \in \mathbb{R}^{3 \times 3}$ uppertriangular. Then $\|\mathbf{B}\mathbf{d}\|$ assumes its minimum on \mathcal{H} in $\tilde{\mathbf{d}}$, where $\tilde{\mathbf{d}}$ is an eigenvector corresponding with the unique positive eigenvector of $\mathbf{R}^{-1}\mathbf{H}\mathbf{R}$.

PROOF. The existence of a minimum follows from compactness and continuity arguments. $(\mathbf{B}^T \mathbf{B} - \lambda \mathbf{H})\mathbf{d} = \mathbf{0}$ implies $(\mathbf{R}^{-T} \mathbf{H} \mathbf{R}^{-1} - \rho^{-1} \mathbf{I})\mathbf{R}\mathbf{d} = \mathbf{0}$. The eigenvalue ρ^{-1} corresponding with the minimum is positive on base of convexity arguments. The positive eigenvalue of $\mathbf{R}^{-T} \mathbf{H} \mathbf{R}^{-1}$ is unique. \square

THEOREM 2.7. Let $\text{rank}(\mathbf{B}) = 1$ and $\text{range}(\mathbf{B}^T) = \text{span}(-\sigma, \mu, \nu)^T$. Then

$$(i) \quad \min\{\|\mathbf{B}\mathbf{d}\| \mid \mathbf{d} \in \mathcal{H}\} = 0, \quad \nu^2 + 4\sigma\mu > 0 \text{ or } \nu = \sigma = \mu = 0;$$

$$(ii) \quad \min\{\|\mathbf{B}\mathbf{d}\| \mid \mathbf{d} \in \mathcal{H}\} = |\nu^2 + 4\sigma\mu|, \quad \nu^2 + 4\sigma\mu < 0;$$

$$(iii) \quad \inf\{\|\mathbf{B}\mathbf{d}\| \mid \mathbf{d} \in \mathcal{H}\} = 0, \quad \nu^2 + 4\sigma\mu = 0 \wedge |\sigma| + |\mu| \neq 0,$$

in this case the infimum is not assumed.

PROOF. In A each block $A_{\ell,m}$ gives a similar contribution to Bd ; hence Bd can be considered to come from one block, say $C = \begin{bmatrix} \alpha & \mu \\ \sigma & \beta \end{bmatrix}$.

- (i) C is diagonalizable with a real shear. The equation $-\sigma d_1 + \mu d_2 + \nu d_3 = 0$ determines a solution curve Γ in \mathcal{H} . The parametric form of Γ is

$$p(\tau) = (\tau, \tau^{-1}, 0)^T, \quad \sigma = \mu = 0 \quad (p_1 > 0)$$

$$p(\tau) = (\nu\tau/\sigma, \sigma(1+\tau^2)(\nu\tau)^{-1}, \tau)^T, \quad \sigma\nu \neq 0, \mu = 0 \quad (p_1 > 0)$$

$$p(\tau) = (-\mu(1+\tau^2)(\nu\tau)^{-1}, -\nu\tau/\mu, \tau)^T, \quad \mu\nu \neq 0, \lambda = 0 \quad (p_1 > 0)$$

$$p(\tau) = \left(\frac{\nu\tau \pm D}{2\sigma}, \frac{-\nu\tau \mp D}{2\mu}, \tau \right)^T, \quad D^2 = (\nu^2 + 4\sigma\mu)\tau^2 + 4\sigma\mu > 0, \lambda\mu \neq 0.$$

- (ii) Transform C in a Murnaghan form. The positive minimum is assumed for

$$d = -|4\sigma\mu + \nu^2|^{\frac{1}{2}}\sigma/|\sigma|(-2\sigma, 2\mu, -\nu)^T.$$

- (iii) C is not diagonalizable. $B^T B$ has no eigenvector in \mathcal{H} . The plane $-\sigma d_1 + \mu d_2 + \nu d_3 = 0$ contacts the recession cone $\{d | d_1 d_2 - d_3^2 = 0, d_1 > 0\}$ of \mathcal{H} along the line $\mathcal{L} : p(\tau) = \tau(2\mu, -2\sigma, -\nu)^T$. Hence $|Bd| > 0$ on \mathcal{H} , for $\mathcal{L} \cap \mathcal{H} = \emptyset$. Now we describe a curve Γ on \mathcal{H} such that \mathcal{L} is its asymptote:

$$\Gamma : d(\tau) = \frac{1}{2}\tau(\mu-\sigma)^{-1}(2\mu, -2\sigma, -\nu) + \frac{1}{2}\tau^{-1}(\mu-\sigma)^{-1}(-2\sigma, 2\mu, \nu), \quad \tau > 0. \quad (2.16)$$

On Γ we find, using the fact that $\nu^2 + 4\sigma\mu = 0$

$$-\sigma d_1(\tau) + \mu d_2(\tau) + \nu d_3(\tau) = (\mu-\sigma)/\tau \rightarrow 0, \quad \tau \rightarrow \infty.$$

The infimum zero of $|Bd|$ on \mathcal{H} is not assumed. \square

EXAMPLE 2.1.

$$A = \begin{bmatrix} -1 & 1 & -2 & 2 \\ -1 & 1 & -2 & 2 \\ -3 & 3 & -4 & 4 \\ -3 & 3 & -4 & 4 \end{bmatrix}$$

Then

$$B^T = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ -2 & -4 & -6 & -8 \end{bmatrix} \text{ and } H^{-1}B^TB = 60 \begin{bmatrix} 1 & 1 & -2 \\ 1 & 1 & -2 \\ 1 & 1 & -2 \end{bmatrix}$$

Remark that $\text{tr}(H^{-1}B^TB) = -(4\sigma\mu + \nu^2) = 0$.

The threefold eigenvalue 0 of nondiagonalizable $H^{-1}B^TB$ gives eigenvectors $(-\gamma_1 + 2\gamma_2, \gamma_1, \gamma_2)^T$, none in \mathcal{H} , for $(-\gamma_1 + 2\gamma_2)\gamma_1 - \gamma_2^2 = -(\gamma_1 - \gamma_2)^2 \leq 0$. With (2.16) we get, since $\sigma = -1$, $\mu = 1$, $\nu = -2$.

$$\Gamma : d(\tau) = \frac{1}{2}(\tau + \tau^{-1}, \tau + \tau^{-1}, \tau - \tau^{-1}) \in \mathcal{H}, \tau > 0.$$

Along Γ holds $\|Bd(\tau)\| \rightarrow 0$ ($\tau \rightarrow 0$). \square

In a similar way one derives the next theorem for the case $\text{rank}(B) = 2$.

THEOREM 2.8. Let be $\text{rank}(B) = 2$ and $\text{nullity}(B) = \text{span}(t_1, t_2, t_3)^T$. Then

- (i) $\min\{\|Bd\| \mid d \in \mathcal{H}\} = 0$, $t_1 t_2 > t_3^2$;
- (ii) $\min\{\|Bd\| \mid d \in \mathcal{H}\} > 0$, $t_1 t_2 < t_3^2$;
- (iii) $\inf\{\|Bd\| \mid d \in \mathcal{H}\} = 0$, $t_1 t_2 = t_3^2$. This infimum is not assumed for finite $d \in \mathcal{H}$. \square

REMARK. In case (iii) $H^{-1}B^TB$ is not diagonalizable. The algebraic multiplicity of eigenvalue zero of $H^{-1}B^TB$ equals two as can be seen from the

coefficient of the first grade term in the characteristic polynomial of $H^{-1}B^TB$, being

$$(t_3^2 - t_1 t_2) \left[\left(\sum_{i=1}^k \mu_i \nu_i \right)^2 - \sum_{i=1}^k \mu_i \sum_{i=1}^k \nu_i \right] / t_1^2 = 0. \quad \square$$

EXAMPLE 2.2.

$$A = \begin{bmatrix} 2 & -2 & 5 & -4 \\ -1 & 1 & -2 & 3 \\ 4 & -4 & 9 & -8 \\ -3 & 0 & -6 & 1 \end{bmatrix}$$

Then

$$B^T = \begin{bmatrix} 1 & 2 & 3 & 6 \\ -2 & -4 & -4 & -8 \\ 1 & 2 & 4 & 8 \end{bmatrix}; \text{rank}(B) = 2, \quad H^{-1}B^TB = \begin{bmatrix} -140 & 200 & -180 \\ 100 & -140 & 130 \\ -65 & 90 & -85 \end{bmatrix}.$$

$\mathcal{N}(B) = \text{span}(4, 1, -2)^T$. Along curve Γ in \mathcal{H} ,

$$\Gamma : \mathbf{d}(\tau) = \frac{1}{5} \tau (3 + 5\sqrt{1+4\tau^{-2}/25}, -3+5\sqrt{1+4\tau^{-2}/25}, -4\tau)^T, \quad \tau > 0$$

holds: $\lim_{\tau \rightarrow \infty} \tau^{-1} \mathbf{d}(\tau) = (4, 1, -2)^T$ and

$$\|\mathbf{B}\mathbf{d}(\tau)\|^2 = \frac{32}{5} (1 + \sqrt{1+4\tau^{-2}/25})^{-2} \tau^{-4} \rightarrow 0 \quad (\tau \rightarrow \infty).$$

$\mathcal{N}(B)$ is the eigenspace of the eigenvalue zero of $H^{-1}B^TB$. \square

3. PARALLEL NORMREDUCTION: COMPLEX MATRICES

Let be A a complex matrix of even order $n = 2k$. The partitioning of A and the notations are as in (2.1) until (2.4). Now we denote

$$x_j = |p_j|^2 + |q_j|^2, \quad y_j = |r_j|^2 + |s_j|^2, \quad z_j = p_j \bar{r}_j + q_j \bar{s}_j, \quad j = 1, \dots, k, \quad (3.1)$$

where p_j, q_j, r_j, s_j are the Jacobi parameters of the complex shear S_j in W . As a consequence of the unimodularity of S_j :

$$x_j y_j - |z_j|^2 = 1, \quad j = 1, \dots, k. \quad (3.2)$$

By simple calculations one derives the following theorem.

THEOREM 3.1. Let be $A' = W^{-1}AW$, $W = \text{diag}(S_1, \dots, S_k)$. Then $\|A'\|_F^2$ is a quadratic function of x_j, y_j, z_j and \bar{z}_j , namely

$$\|A'\|_F^2 = \sum_{\ell, m=1}^k (x_\ell, y_\ell, z_\ell, \bar{z}_\ell) B_{\ell, m} (x_m, y_m, z_m, \bar{z}_m)^T =$$

where

$$B_{\ell, m} = \begin{bmatrix} |\sigma_{\ell, m}|^2 & |\beta_{\ell, m}|^2 & \sigma_{\ell, m} \bar{\beta}_{\ell, m} & \bar{\sigma}_{\ell, m} \beta_{\ell, m} \\ |\alpha_{\ell, m}|^2 & |\mu_{\ell, m}|^2 & \alpha_{\ell, m} \bar{\mu}_{\ell, m} & \bar{\alpha}_{\ell, m} \mu_{\ell, m} \\ -\bar{\alpha}_{\ell, m} \sigma_{\ell, m} & -\bar{\mu}_{\ell, m} \beta_{\ell, m} & -\sigma_{\ell, m} \bar{\mu}_{\ell, m} & -\bar{\alpha}_{\ell, m} \beta_{\ell, m} \\ -\alpha_{\ell, m} \bar{\sigma}_{\ell, m} & -\mu_{\ell, m} \bar{\beta}_{\ell, m} & -\alpha_{\ell, m} \beta_{\ell, m} & -\bar{\sigma}_{\ell, m} \mu_{\ell, m} \end{bmatrix}. \quad (3.3)$$

□

Let be

$$x_j = t_j + w_j, \quad y_j = t_j - w_j, \quad j = 1, \dots, k \quad (3.4)$$

Then, as follows from the unimodularity of S_j ,

$$t_j = (1 + w_j^2 + z_j \bar{z}_j)^{\frac{1}{2}}.$$

Hence $\|A'\|_F^2$ is a function of the k triples (w_j, z_j, \bar{z}_j) , $j = 1, \dots, k$:

$$\|A'\|_F^2 = g(w_1, z_1, \bar{z}_1, \dots, w_k, z_k, \bar{z}_k).$$

The relation between the commutator $C^{(')} = A^{(')}^* A^{(')} - A^{(')} A^{(')}^*$ and g is mentioned in

THEOREM 3.2. The partial derivatives of g in $0 = (0, 0, 0, \dots, 0, 0, 0) \in \mathbb{C}^{3k}$ satisfy the following densities

$$\begin{cases} \frac{\partial g}{\partial w_j}(0) = c_{2j-1, 2j-1} - c_{2j, 2j}, \\ \frac{\partial g}{\partial z_j}(0) = c_{2j, 2j-1}, \\ \frac{\partial g}{\partial \bar{z}_j}(0) = c_{2j-1, 2j}. \end{cases} \quad j = 1, \dots, k \quad (3.5)$$

PROOF. Use the properties

$$\frac{\partial x_j}{\partial w_j} = 1 + w_j/t_j = x_j/t_j, \quad \frac{\partial y_j}{\partial w_j} = 1 - w_j/t_j = y_j/t_j, \quad \frac{\partial x_j}{\partial z_j} = \bar{z}_j/t_j$$

and

$$c_{2j-1, 2j-1} - c_{2j, 2j} = \sum_{m=1}^k (|\alpha_{m,j}|^2 + |\lambda_{m,j}|^2 - |\mu_{m,j}|^2 - |\beta_{m,j}|^2 + |\lambda_{j,m}|^2 + |\beta_{j,m}|^2 - |\alpha_{j,m}|^2 - |\mu_{j,m}|^2),$$

$$c_{2j-1, 2j} = \sum_{m=1}^k (\bar{\alpha}_{m,j} \mu_{m,j} + \bar{\lambda}_{m,j} \beta_{m,j} - \alpha_{j,m} \bar{\lambda}_{j,m} - \mu_{j,m} \bar{\beta}_{j,m}).$$

With (3.3), and (3.4) and easy but cumbersome calculation one finds (3.5). \square

As in the preceding section we restrict ourselves to a *diagonal of shears*:

$$W = \text{diag}(S_1, \dots, S_k) \quad (3.6)$$

with

$$S_j = S = \begin{pmatrix} p & q \\ r & s \end{pmatrix}, \quad ps - qr = 1, \quad j = 1, \dots, k. \quad (3.7)$$

Similar to (2.8) we define common Euclidean parameters

$$x = |p|^2 + |q|^2, \quad y = |r|^2 + |s|^2, \quad z = p\bar{r} + q\bar{s} = u + iv \quad (3.8)$$

with

$$xy - |z|^2 = 1, \quad (3.9)$$

where $|z|^2 = u^2 + v^2$. Further

$$\mathcal{H} = \{(x, y, u, v) \mid x > 0, \quad xy - u^2 - v^2 = 1\}.$$

By rather simple calculations $\|W^{-1}AW\|_F^2$ appears to be a quadratic function of x, y, u and v .

THEOREM 3.3. If W is a diagonal of shears with common Euclidean parameters $x, y, z = u + iv$ then $\|W^{-1}AW\|_F^2$ is expressible in terms of these parameters, viz.

$$\begin{aligned} \|W^{-1}AW\|_F^2 &= f(x, y, z) = \sum_{\ell, m=1}^k (|\alpha_{\ell, m}|^2 + |\beta_{\ell, m}|^2 + |\mu_{\ell, m}y + \nu_{\ell, m}z|^2 \\ &\quad + |\sigma_{\ell, m}x - \nu_{\ell, m}\bar{z}|^2 - |\nu_{\ell, m}z|^2 - 2 \operatorname{Re}(\bar{\sigma}_{\ell, m}\mu_{\ell, m}z^2)) = \\ &= \sum_{\ell, m=1}^k (|\alpha_{\ell, m}|^2 + |\beta_{\ell, m}|^2) + (x, y, u, v) \sum_{\ell, m=1}^k P_{\ell, m}(x, y, u, v)^T, \end{aligned} \quad (3.10)$$

where

$$\begin{bmatrix} |\sigma_{\ell,m}|^2 & 0 & -\operatorname{Re}(\bar{\sigma}_{\ell,m} \nu_{\ell,m}) & -\operatorname{Im}(\bar{\sigma}_{\ell,m} \nu_{\ell,m}) \\ 0 & |\mu_{\ell,m}|^2 & \operatorname{Re}(\mu_{\ell,m} \bar{\nu}_{\ell,m}) & \operatorname{Im}(\mu_{\ell,m} \bar{\nu}_{\ell,m}) \\ -\operatorname{Re}(\bar{\sigma}_{\ell,m} \nu_{\ell,m}) & \operatorname{Re}(\mu_{\ell,m} \bar{\nu}_{\ell,m}) & |\nu_{\ell,m}|^2 - 2\operatorname{Re}(\sigma_{\ell,m} \bar{\mu}_{\ell,m}) & 2\operatorname{Im}(\sigma_{\ell,m} \bar{\mu}_{\ell,m}) \\ -\operatorname{Im}(\bar{\sigma}_{\ell,m} \nu_{\ell,m}) & \operatorname{Im}(\mu_{\ell,m} \bar{\nu}_{\ell,m}) & 2\operatorname{Im}(\sigma_{\ell,m} \bar{\mu}_{\ell,m}) & |\nu_{\ell,m}|^2 + 2\operatorname{Re}(\sigma_{\ell,m} \bar{\mu}_{\ell,m}) \end{bmatrix} \quad \square \quad (3.11)$$

Analogously to section 2 we define

$$w = (x-g)/2, \quad t = (x+y)/2 = \sqrt{1+u^2+v^2+w^2}. \quad (3.12)$$

Then, as follows from (3.10), $\|W^{-1}AW\|_F^2$ is a function of w , u and v :

$$g(w,u,v;A) := \|W^{-1}AW\|_F^2. \quad (3.13)$$

With simple but cumbersome calculations one proves the following lemma.

LEMMA 3.4. Let be $(c'_{i,j}) = (A')^*A' - A'(A')^*$, where $A' = W^{-1}AW$ with W as defined in (3.6) and (3.7). Then

$$\sum_{\ell=1}^k \begin{bmatrix} c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell} \\ c'_{2\ell-1,2\ell} \\ c'_{2\ell,2\ell-1} \end{bmatrix} = \frac{1}{2} \begin{bmatrix} |p|^2 + |s|^2 - |q|^2 - |r|^2 & \bar{p}r - \bar{q}s & p\bar{r} - q\bar{s} \\ \bar{p}q - \bar{r}s & \bar{p}s & q\bar{r} \\ p\bar{q} - r\bar{s} & \bar{q}r & p\bar{s} \end{bmatrix} \begin{bmatrix} g_w \\ g_u + ig_v \\ g_u - ig_v \end{bmatrix} \quad (3.14)$$

where g as defined in (3.11). \square

THEOREM 3.5. The function g is stationary in $(w,u,v) \in \mathbb{R}^3$ iff

$$\sum_{\ell=1}^k (c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell}) = 0 \quad \text{and} \quad \sum_{\ell=1}^k c'_{2\ell-1,2\ell} = 0.$$

PROOF. The determinant of the coefficient matrix in (3.12) equals $\frac{1}{2}|ps - qr|^2(|p|^2 + |q|^2 + |r|^2 + |s|^2) = t > 0$. This proves the theorem. \square

Theorem 3.3 implies that the determination of the optimal normreducing diagonal of shears requires the minimization of a quadratic function subject to $xy - u^2 - v^2 = 1$. Neither the function f in (3.10) nor $h(x, y, z) = xy - |z|^2$ are analytic, in contrast with the corresponding real functions (2.10) and (2.9) resp.

Let be $\mathbf{d} = (d_1, d_2, d_3, d_4) = (x, y, u, v)$,

$$H = \begin{bmatrix} 0 & \frac{1}{2} & 0 & 0 \\ \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \quad (3.15)$$

and

$$\mathcal{H} = \{\mathbf{d} \in \mathbb{R}^4 \mid \mathbf{d}^T H \mathbf{d} = 1, d_1 > 0\} \quad (3.16)$$

Further $M = \sum_{\ell, m=1}^k P_{\ell, m}$, where $P_{\ell, m} \in \mathbb{R}^{4 \times 4}$ defined in theorem 3.3 and let be $h(\mathbf{d}) := \mathbf{d}^T M \mathbf{d}$, $\mathbf{d} \in \mathbb{R}^4$.

With these notations the minimization problem becomes: solve

$$\min\{h(\mathbf{d}) \mid \mathbf{d} \in \mathcal{H}\}. \quad (3.17)$$

Let be λ_j , $j = 1, \dots, n$ the eigenvalues of A . Then

$$\sum_{j=1}^n |\lambda_j|^2 \leq \|W^{-1} A W\|_F^2 = h(\mathbf{d}) + \sum_{\ell, m=1}^k (|\alpha_{\ell, m}|^2 + |\beta_{\ell, m}|^2)$$

for each $\mathbf{d} \in \mathcal{H}$. So h is bounded below on \mathcal{H} . Let be

$$\eta = \inf\{h(\mathbf{d}) \mid \mathbf{d} \in \mathcal{H}\}. \quad (3.18)$$

THEOREM 3.6. Let be η as defined in (3.18).

(i) If infimum η is assumed in $\mathbf{d} \in \mathcal{H}$ then $M \mathbf{d} = \eta H \mathbf{d}$.

(ii) If $h(\mathbf{x}) > \eta$ for each $\mathbf{x} \in \mathcal{H}$ then there exist a $\mathbf{d} = (d_1, d_2, d_3, d_4)^T$ such that $M \mathbf{d} = \eta H \mathbf{d}$, $\|\mathbf{d}\| = 1$, $d_1 > 0$ and $\mathbf{d}^T H \mathbf{d} = 0$.

PROOF.

- (i) The Lagrange multiplier method gives $Md = \lambda Hd$ and $\eta = d^T Md = \lambda d^T Hd = \lambda$. So $Md = \eta Hd$.
- (ii) Let be $\mathcal{K} = \{d \in \mathbb{R}^4 | d^T Hd \geq 0\}$ and $S = \{x \in \mathcal{K} | \|x\| = 1\}$. Without loss of generality we may assume $\eta = 0$. There exists a sequence $\{x_n\}$ in \mathcal{K} and a corresponding sequence $\{\hat{x}_n\}$ in S , with $\hat{x}_n = x_n / \|x_n\|$, such that

$$x_n^T M x_n = \hat{x}_n^T M x_n / \hat{x}_n^T H \hat{x}_n \downarrow \eta \quad (, n \rightarrow \infty) . \quad (3.19)$$

Some subsequence $\{\hat{x}_{n_k}\}$ of $\{\hat{x}_n\}$ is convergent; let be d its limit; $d \in \partial \mathcal{K}$ for otherwise the infimum η would be assumed on \mathcal{K} . So $d^T Hd = 0$. Hence also $d^T Md = 0$. It is clear that $x^T M x \geq 0$ for each $x \in \mathcal{K}$. So $h|_{\partial \mathcal{K}}$ assumed its minimum in d . Application of Lagranges multiplier method for that minimum gives: there exists a $\lambda \in \mathbb{R}$ such that

$$Md = \lambda Hd . \quad (3.20)$$

Let be $x = d + h \in S$. Since $d^T Md = d^T Hd$ we get with (3.20)

$$\frac{x^T M x}{x^T H x} = \frac{2\lambda(Bd)^T h + h^T M h}{2(Bd)^T h + h^T H h} . \quad (3.21)$$

Now $(Bd)^T h > 0$ when $h \in \text{int}(\mathcal{K})$. Hence

$$x^T M x / x^T H x \rightarrow \lambda \quad (, x \in S, x \rightarrow d) .$$

So $\lambda = \eta$. \square

COROLLARY. If $x^T M x > \eta$ for each $x \in \mathcal{K}$ then the intersection of \mathcal{K} and the subspace $\{\tau_1(1,1,0,0)^T + \tau_2 d | \tau_1, \tau_2 > 0\}$ is a curve

$$\Gamma : x(t) = t(d_1 + d_2)^{-1} d + t^{-1}(d_1 + d_2)^{-1}(d_2, d_1, -d_3, -d_4)^T$$

along which $x(t)^T M x(t)$ tends to η for $t \rightarrow \infty$. \square

EXAMPLE 3.1.

Let be

$$A = \begin{bmatrix} 0 & -i & 1 & 2 \\ -i & 2 & 2 & 1+4i \\ 4+i & 1-3i & 5+i & 2+i \\ 1-3i & 10+3i & 2+i & 3+5i \end{bmatrix}$$

Then

$$M = \begin{bmatrix} 20 & 0 & 0 & 40 \\ 0 & 20 & 0 & 40 \\ 0 & 0 & 40 & 0 \\ 40 & 40 & 0 & 120 \end{bmatrix}, \quad H^{-1}M = 40 \begin{bmatrix} 0 & 1 & 0 & 2 \\ 1 & 0 & 0 & 2 \\ 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & -3 \end{bmatrix}.$$

$\lambda = -40$ is the fourfold eigenvalue of the pair (M, H) . The eigenspace is spanned by the three vectors $(-1, 1, 0, 0)^T$, $(-2, 0, 0, 1)^T$ and $(0, 0, 1, 0)^T$. None linear combination of these vectors is in \mathcal{H} ; $d = (\sqrt{3})^{-1}(1, 1, 0, -1)^T$ satisfies the equality $d_1 d_2 - d_3^2 - d_4^2 = 0$. The line $x(\tau) = \tau(1, 1, 0, -1)^T$ is the asymptote of the curve $\Gamma : x(\tau) = (2\tau)^{-1}(\tau^2+1, \tau^2+1, 0, \tau^2-1)^T$ in \mathcal{H} . Γ is the intersection of the plane $\{\tau_1(1, 1, 0, 0)^T + \tau_2 d \mid \tau_1, \tau_2 > 0\}$ and \mathcal{H} . Along Γ holds $h(x(\tau)) \rightarrow \eta = -40$. Remark that $-40 + \sum_{\lambda, m=1}^4 (|\alpha_{\lambda, m}|^2 + |\beta_{\lambda, m}|^2) = 168$, being the sum of the squares of the moduli of the eigenvalues of the four non-diagonalizable matrices $A_{\lambda, m}$ of A . \square

4. PRECONDITIONED PARALLEL NORMREDUCTION

This section describes a preconditioning transformation by unitary matrices, and the following suboptimal normreduction. This combination brings about convergence to normality. In the first step that twofold action results in

$$A' = W^{-1}U^*AUW = W^{-1}(U^*AU)W \quad (4.1)$$

where

$$U = \text{diag}(U_1, \dots, U_k), \quad W = \text{diag}(S, \dots, S) \quad (4.2)$$

with

$$U_\ell = \begin{bmatrix} \cos \varphi_\ell & -e^{-i\vartheta_\ell} \sin \varphi_\ell \\ e^{i\vartheta_\ell} \sin \varphi_\ell & \cos \varphi_\ell \end{bmatrix}, \quad \ell = 1, \dots, k \quad (4.3)$$

and

$$S = \begin{bmatrix} p & 0 \\ 0 & s \end{bmatrix}, \quad ps = 1$$

a 2×2 submatrix in the diagonal of shears W .

As in Sameh's algorithm [5] the *rectifier* U is chosen such that $\text{grad } g(0,0,0; W^{-1}AW)$ (see (2.11) and (3.17)) has maximal length. For complex A we use theorem 3.4.

THEOREM 4.1. Let be $U = \text{diag}(U_1, \dots, U_k)$, with U_ℓ , $\ell = 1, \dots, k$, as given in (4.3) and let be

$$v_\ell(A) = v_\ell = \begin{bmatrix} c_{2\ell-1, 2\ell-1} - c_{2\ell, 2\ell} \\ 2c_{2\ell-1, 2\ell} \end{bmatrix}, \quad \ell = 1, \dots, k \quad (4.4)$$

where $(c_{i,j}) = C(A) = A^*A - AA^*$. Then the maximum length of $\text{grad } g(0,0,0;U^*AU)$ with respect to U equals $\sum_{\ell=1}^k |v_\ell|$ and is assumed for

$$\begin{bmatrix} \cos 2\varphi_\ell \\ \sin 2\varphi_\ell \end{bmatrix} = \begin{cases} (1,0)^T, & \text{if } v_\ell = 0, \\ (c_{2\ell-1,2\ell-1} - c_{2\ell,2\ell}, 2(c_{2\ell-1,2\ell})) |v_\ell|^{-1}, & \text{if } v_\ell \neq 0 \end{cases} \quad (4.5)$$

with $|\varphi_\ell| \leq \frac{\pi}{2}$ and

$$e^{i\vartheta} = \begin{cases} 1, & \text{if } c_{2\ell-1,2\ell} = 0 \\ (c_{2\ell,2\ell-1}/|c_{2\ell-1,2\ell}|, \text{ if } c_{2\ell-1,2\ell} \neq 0. \end{cases} \quad (4.6)$$

Then $g_w(0,0,0;U^*AU) \geq 0$, $g_u(0,0,0;U^*AU) = g_v(0,0,0;U^*AU) = 0$.

PROOF. It follows from the corollary of theorem 3.4 that $\text{grad } g(0,0,0;U^*AU) = \sum_{\ell=1}^k (c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell}, 2\text{Re}(c_{2\ell-1,2\ell}), 2\text{Im}(c_{2\ell-1,2\ell}))$ where $(c'_{i,j}) = C(U^*AU)$. Since $C(U^*AU) = U^*C(A)U$ we find

$$\begin{bmatrix} c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell} \\ e^{i\vartheta} c'_{2\ell-1,2\ell} \end{bmatrix} = \begin{bmatrix} \cos 2\varphi & \sin 2\varphi & \sin 2\varphi \\ -\frac{1}{2}\sin 2\varphi & \cos^2 \varphi & -\sin^2 \varphi \end{bmatrix} \begin{bmatrix} c_{2\ell-1,2\ell-1} - c_{2\ell,2\ell} \\ e^{i\vartheta} c_{2\ell-1,2\ell} \\ e^{-i\vartheta} \bar{c}_{2\ell-1,2\ell} \end{bmatrix}$$

With $\cos \varphi$, $\sin \varphi$ and $e^{i\vartheta}$ as given in (4.6) and (4.5)

$$\begin{bmatrix} c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell} \\ 2c_{2\ell-1,2\ell} \end{bmatrix} = v_\ell(U^*AU) = \begin{bmatrix} |v_\ell(A)| \\ 0 \end{bmatrix}. \quad (4.7)$$

With these transformations U_1, \dots, U_k the k vectors $v_\ell(U^*AU)$ have the same direction, the vectors $v_\ell(A)$ are *rectified*. Thus

$$|g(0,0,0;U^*AU)| = \left| \sum_{\ell=1}^k v_\ell(U^*AU) \right| = \sum_{\ell=1}^k |v_\ell(U^*AU)| = \sum_{\ell=1}^k |v_\ell(A)|.$$

Now $g_u(0,0,0;U^*AU) = g_v(0,0,0;U^*AU) = 0$ and $g_w(0,0,0;U^*AU) \geq 0$. \square

THEOREM 4.2. Let be U a unitary matrix as defined in (4.2), (4.3), (4.5) and (4.6). Then there exists a diagonal matrix W such that

$$\|A\|_F^2 - \|W^{-1}U^*AUW\|_F^2 \geq \frac{1}{8}\|A\|_F^2 \sum_{\ell=1}^k |v_\ell|^2 \quad (4.8)$$

where v_ℓ as defined in (4.4).

PROOF. Let be $A' = U^*AU$ and W a diagonal of identical diagonal shears $\begin{pmatrix} p & 0 \\ 0 & s \end{pmatrix}$. According to (3.12)

$$\|W^{-1}A'W\|_F^2 = \sum_{\ell,m=1}^k (|\alpha'_{\ell,m}|^2 + |\beta'_{\ell,m}|^2 + |\mu'_{\ell,m}|^2 x^{-2} + |\sigma'_{\ell,m}|^2 x^2),$$

for the Euclidean parameter z of a diagonal shear equals zero. Let be $\sum_{\ell,m=1}^k |\sigma'_{\ell,m}|^2 = c_1$, $\sum_{\ell,m=1}^k |\mu'_{\ell,m}|^2 = c_2$.

1. Let be $c_1 c_2 \neq 0$. Then $c_1 x^2 + c_2 x^{-2}$ is minimal for $x = (c_2/c_1)^{\frac{1}{4}}$. With the Euclidean parameters $((c_2/c_1)^{\frac{1}{4}}, (c_1/c_2)^{\frac{1}{4}}, 0)$ the decrease of the Euclidean norm equals

$$\|A\|_F^2 - \|W^{-1}U^*AUW\|_F^2 = (\sqrt{c_1} - \sqrt{c_2})^2.$$

Since $c_1 - c_2 = \frac{1}{2} g_w(0,0,0;A')$

$$\sqrt{c_1} - \sqrt{c_2} = (\sqrt{c_1} + \sqrt{c_2})^{-1} (c_1 - c_2) = \frac{1}{2} (\sqrt{c_1} + \sqrt{c_2})^{-1} g_w(0,0,0;A').$$

Now $\sqrt{c_1} + \sqrt{c_2} \leq \sqrt{2}(c_1 + c_2)^{\frac{1}{2}} \leq \sqrt{2}\|A'\|_F = \sqrt{2}\|A\|_F$ and

$$g_w(0,0,0;A') = \sum_{\ell=1}^k (c'_{2\ell-1,2\ell-1} - c'_{2\ell,2\ell}) = \sum_{\ell=1}^k |v_\ell(U^*AU)| \geq \sum_{\ell=1}^k |v_\ell(A)|$$

as can be seen from (4.7). Consequently

$$\|A\|_F^2 - \|W^{-1}U^*AUW\|_F^2 \geq \frac{1}{8} \|A\|_F^{-2} \left(\sum_{\ell=1}^k \|v_\ell(A)\| \right)^2 \geq \frac{1}{8} \|A\|_F^{-2} \sum_{\ell=1}^k \|v_\ell(A)\|^2.$$

2. In case $c_1 \neq 0$ and $c_2 = 0$, choose x so small that $x^2 \leq 1 - \|A\|_F^{-2} c_1/8$. Then

$$\|A\|_F^2 - \|W^{-1}U^*AUW\|_F^2 = c_1(1-x^2) \geq \|A\|_F^{-2} c_1^2/8 \geq \frac{1}{8} \|A\|_F^{-2} \sum_{\ell=1}^k \|v_\ell(A)\|^2$$

$$\text{for } c_1 = \sum_{\ell=1}^k (c'_{2\ell-1, 2\ell-1} - c'_{2\ell, 2\ell}) = \sum_{\ell=1}^k \|v_\ell(U^*AU)\| = \sum_{\ell=1}^k \|v_\ell(A)\|.$$

3. In case $c_2 = 0$ and $c_1 \neq 0$ choose $x^2 \geq (1 - \|A\|_F^{-2} c_2/8)^{-1}$. \square

The pivot strategy guaranteeing that the Euclidean norm decreases in sufficient degree for convergence to normality [2,3] will be derived from lower bound (4.8). Therefore we need

THEOREM 4.3. There exists a set of k distinct index pairs (ℓ_j, m_j) , with $\ell_j < m_j$, $j = 1, \dots, k$, such that

$$\sum_{j=1}^k (c_{\ell_j, \ell_j} - c_{m_j, m_j}) + 4|c_{\ell_j, m_j}|^2 \geq \frac{4}{n-1} \|C(A)\|_F^2. \quad (4.9)$$

PROOF. We have $\sum_{\ell \neq m} (c_{\ell, \ell} - c_{m, m})^2 = 2(n-1) \sum_{\ell=1}^n c_{\ell, \ell}^2 - 2 \sum_{\ell \neq m} c_{\ell, \ell} c_{m, m}$. But since $\sum_{\ell=1}^n c_{\ell, \ell} = 0$, $(\sum_{\ell=1}^n c_{\ell, \ell})^2 = \sum_{\ell=1}^n c_{\ell, \ell}^2 + \sum_{\ell \neq m} c_{\ell, \ell} c_{m, m} = 0$. Hence for $n \geq 2$: $\sum_{\ell \neq m} (c_{\ell, \ell} - c_{m, m})^2 = 2n \sum_{\ell=1}^n c_{\ell, \ell}^2 \geq 4 \sum_{\ell=1}^n c_{\ell, \ell}^2$. Consequently

$$\sum_{\ell \neq m} ((c_{\ell, \ell} - c_{m, m})^2 + 4|c_{\ell, m}|^2) \geq 4\|C(A)\|_E^2 \quad (4.10)$$

Let be \mathcal{Q} the collection of all sets ω of k distinct index pairs (ℓ_j, m_j) . The number of sets ω in \mathcal{Q} is $n!/(k!2^k)$, and each pair (ℓ, m) , with $\ell \neq m$, occurs in $(n-2)!/((k-1)!2^{k-1})$ sets of \mathcal{Q} . Thus

$$\begin{aligned} \sum_{\omega \in \Omega} \sum_{(\ell, m) \in \omega} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2) &= \sum_{\ell \neq m} \sum_{\omega \in \Omega} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2) \\ &= \frac{(n-2)!}{(k-1)!2^{k-1}} \sum_{\ell \neq m} \sum_{(\ell, m) \in \omega} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2) . \end{aligned}$$

Hence the mean of $\sum_{(\ell, m) \in \omega} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2)$ overall $\omega \in \Omega$ equals

$$\begin{aligned} \left[\frac{n!}{k!2^k} \right]^{-1} \frac{(n-2)!}{(k-1)!2^{k-1}} \sum_{\ell \neq m} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2) &= \\ (n-1)^{-1} \sum_{\ell \neq m} ((c_{\ell, \ell^{-c_{m, m}}})^2 + 4|c_{\ell, m}|^2) . \end{aligned}$$

This result, together with (4.10), proves the theorem. \square

THEOREM 4.4. Let a sequence $\{A^{(j)}\}$, starting with $A^{(0)} = A$, be constructed by

$$A^{(j+1)} = (P^{(j)} U^{(j)} W^{(j)})^{-1} A^{(j)} P^{(j)} U^{(j)} W^{(j)} , \quad j = 1, 2, \dots \quad (4.11)$$

where in each step k disjunct index pairs $(\ell_{1,j}, m_{1,j}), \dots, (\ell_{k,j}, m_{k,j})$ are selected according to rule (4.9). $P^{(j)}$ is a permutation with $P(\ell_{1,j}, m_{1,j}, \dots, \ell_{k,j}, m_{k,j}) = (1, 2, \dots, n-1, n)$. $U^{(j)}$ is a preconditioning unitary block diagonal matrix as described in (4.2), (4.3), (4.5) and (4.6) and $\tilde{W}^{(j)} = \text{diag}(x_j^{\frac{1}{2}}, x_j^{-\frac{1}{2}}, \dots, x_j^{\frac{1}{2}}, x_j^{-\frac{1}{2}})$ that reduces the Frobenius norm of $(P^{(j)} U^{(j)})^{-1} A^{(j)} P^{(j)} U^{(j)}$ as described in theorem 4.2. Then $\{A^{(j)}\}$ converges to normality.

PROOF. $\{A_F^{(j)}\}$ decreases monotonically and is bounded below. Therefore $\delta_j := \|A^{(j)}\|_F^2 - \|A^{(j+1)}\|_F^2 \downarrow 0, (j \rightarrow \infty)$. Since by theorem 4.2 and theorem 4.3

$$\begin{aligned}
\delta_j &\geq \frac{1}{8} \|A\|_F^{-2} \sum_{k=1}^k \left\| v_k \left((P^{(j)} U^{(j)})^{-1} A^{(j)} P^{(j)} U^{(j)} \right) \right\|_F^2 \\
&\geq \frac{1}{2(n-1)} \|A\|_F^{-2} \|C((P^{(j)} U^{(j)}) A^{(j)} P^{(j)} U^{(j)})\|_F^2 \\
&= \frac{1}{2(n-1)} \|A\|_F^{-2} \|C(A^{(j)})\|_F^2,
\end{aligned}$$

we have $C(A^{(j)}) \rightarrow 0$ ($j \rightarrow \infty$). \square

THEOREM 4.5. Let $\{A^{(j)}\}$ be constructed recursively by

$$A^{(j+1)} = (P^{(j)} U^{(j)} W^{(j)})^{-1} A^{(j)} P^{(j)} U^{(j)} W^{(j)}, \quad j = 1, 2, \dots \quad (4.12)$$

with $P^{(j)}$ and $U^{(j)}$ as in theorem 4.4 but $W^{(j)}$ an optimal parallel norm-reducing shear as described in section 3. Then $\{A^{(j)}\}$ converges to normality.

PROOF. $\delta_j = \|A^{(j)}\|_F^2 - \|A^{(j+1)}\|_F^2 \downarrow 0$ ($j \rightarrow \infty$) for now $W^{(j)}$ is even optimal. As in the preceding theorem $C(A^{(j)}) \rightarrow 0$ ($j \rightarrow \infty$). \square

REFERENCES

1. P.J. Eberlein (1962). A Jacobi-like Method for the Automatic Computation of Eigenvalues and Eigenvectors of an Arbitrary Matrix, J. Soc. Indust. Appl. Math., 10, pp. 74-88.
2. L. Elsner, M.H.C. Paardekooper (1987). On Measures of Nonnormality of Matrices, Lin. Algebra Appl., 92, 107-124.
3. M.H.C. Paardekooper (1969). An Eigenvalue Algorithm Based on Normreducing Transformations, Technological University Eindhoven, 144 pp.
4. M.H.C. Paardekooper (1987). Sameh's Parallel Eigenvalue Algorithm Revisited, in Algorithm and Applications of Vector and Parallel Computers, (H.J.J. te Riele, T.J. Dekker and H.A. van der Vorst, Eds.), Elseviers Science Publishers, pp. 351-371.
5. A.H. Sameh (1971). On Jacobi and Jacobi-like Algorithms for a Parallel Computer, Math. Comp., 25, pp. 579-590.
6. K. Veselić (1976). A Convergence Jacobi Method for Solving the Eigenproblem of Arbitrary Real Matrices, Num. Math., 25, pp. 178-184.

IN 1989 REEDS VERSCHENEN

- 368 Ed Nijssen, Will Reijnders
"Macht als strategisch en tactisch marketinginstrument binnen de distributieketen"
- 369 Raymond Gradus
Optimal dynamic taxation with respect to firms
- 370 Theo Nijman
The optimal choice of controls and pre-experimental observations
- 371 Robert P. Gilles, Pieter H.M. Ruys
Relational constraints in coalition formation
- 372 F.A. van der Duyn Schouten, S.G. Vanneste
Analysis and computation of (n,N)-strategies for maintenance of a two-component system
- 373 Drs. R. Hamers, Drs. P. Verstappen
Het company ranking model: a means for evaluating the competition
- 374 Rommert J. Casimir
Infogame Final Report
- 375 Christian B. Mulder
Efficient and inefficient institutional arrangements between governments and trade unions; an explanation of high unemployment, corporatism and union bashing
- 376 Marno Verbeek
On the estimation of a fixed effects model with selective non-response
- 377 J. Engwerda
Admissible target paths in economic models
- 378 Jack P.C. Kleijnen and Nabil Adams
Pseudorandom number generation on supercomputers
- 379 J.P.C. Blanc
The power-series algorithm applied to the shortest-queue model
- 380 Prof. Dr. Robert Bannink
Management's information needs and the definition of costs, with special regard to the cost of interest
- 381 Bert Bettonvil
Sequential bifurcation: the design of a factor screening method
- 382 Bert Bettonvil
Sequential bifurcation for observations with random errors

- 383 Harold Houba and Hans Kremers
Correction of the material balance equation in dynamic input-output models
- 384 T.M. Doup, A.H. van den Elzen, A.J.J. Talman
Homotopy interpretation of price adjustment processes
- 385 Drs. R.T. Frambach, Prof. Dr. W.H.J. de Freytas
Technologische ontwikkeling en marketing. Een oriënterende beschouwing
- 386 A.L.P.M. Hendrikx, R.M.J. Heuts, L.G. Hoving
Comparison of automatic monitoring systems in automatic forecasting
- 387 Drs. J.G.L.M. Willems
Enkele opmerkingen over het inversificerend gedrag van multinationale ondernemingen
- 388 Jack P.C. Kleijnen and Ben Annink
Pseudorandom number generators revisited
- 389 Dr. G.W.J. Hendrikse
Speltheorie en strategisch management
- 390 Dr. A.W.A. Boot en Dr. M.F.C.M. Wijn
Liquiditeit, insolventie en vermogensstructuur
- 391 Antoon van den Elzen, Gerard van der Laan
Price adjustment in a two-country model
- 392 Martin F.C.M. Wijn, Emanuel J. Bijnen
Prediction of failure in industry
An analysis of income statements
- 393 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters
On the short term objectives of daily intervention by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 394 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters
On the effectiveness of daily interventions by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 395 A.E.M. Meijer and J.W.A. Vingerhoets
Structural adjustment and diversification in mineral exporting developing countries
- 396 R. Gradus
About Tobin's marginal and average q
A Note
- 397 Jacob C. Engwerda
On the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = I$

- 398 Paul C. van Batenburg and J. Kriens
Bayesian discovery sampling: a simple model of Bayesian inference in auditing
- 399 Hans Kremers and Dolf Talman
Solving the nonlinear complementarity problem
- 400 Raymond Gradus
Optimal dynamic taxation, savings and investment
- 401 W.H. Haemers
Regular two-graphs and extensions of partial geometries
- 402 Jack P.C. Kleijnen, Ben Annink
Supercomputers, Monte Carlo simulation and regression analysis
- 403 Ruud T. Frambach, Ed J. Nijssen, William H.J. Freytas
Technologie, Strategisch management en marketing
- 404 Theo Nijman
A natural approach to optimal forecasting in case of preliminary observations
- 405 Harry Barkema
An empirical test of Holmström's principal-agent model that tax and signally hypotheses explicitly into account
- 406 Drs. W.J. van Braband
De begrotingsvoorbereiding bij het Rijk
- 407 Marco Wilke
Societal bargaining and stability
- 408 Willem van Groenendaal and Aart de Zeeuw
Control, coordination and conflict on international commodity markets
- 409 Prof. Dr. W. de Freytas, Drs. L. Arts
Tourism to Curacao: a new deal based on visitors' experiences
- 410 Drs. C.H. Veld
The use of the implied standard deviation as a predictor of future stock price variability: a review of empirical tests
- 411 Drs. J.C. Caanen en Dr. E.N. Kertzman
Inflatieneutrale belastingheffing van ondernemingen
- 412 Prof. Dr. B.B. van der Genugten
A weak law of large numbers for m -dependent random variables with unbounded m
- 413 R.M.J. Heuts, H.P. Seidel, W.J. Selen
A comparison of two lot sizing-sequencing heuristics for the process industry

- 414 C.B. Mulder en A.B.T.M. van Schaik
Een nieuwe kijk op structuurwerkloosheid
- 415 Drs. Ch. Caanen
De hefboomwerking en de vermogens- en voorraadaftrek
- 416 Guido W. Imbens
Duration models with time-varying coefficients
- 417 Guido W. Imbens
Efficient estimation of choice-based sample models with the method of moments
- 418 Harry H. Tigelaar
On monotone linear operators on linear spaces of square matrices

IN 1990 REEDS VERSCHENEN

- 419 Bertrand Melenberg, Rob Alessie
A method to construct moments in the multi-good life cycle consumption model
- 420 J. Kriens
On the differentiability of the set of efficient (μ, σ^2) combinations in the Markowitz portfolio selection method
- 421 Steffen Jørgensen, Peter M. Kort
Optimal dynamic investment policies under concave-convex adjustment costs
- 422 J.P.C. Blanc
Cyclic polling systems: limited service versus Bernoulli schedules

Bibliotheek K. U. Brabant



17 000 01066393 9